arXiv:1402.0995v1 [physics.bio-ph] 5 Feb 2014

The role of carbohydrates at the origin of homochirality in biosystems

Søren Toxvaerd

(Dated: February 6, 2014)

Abstract

Pasteur has demonstrated that the chiral components in a racemic mixture can separate in homochiral crystals. But with a strong chiral discrimination the chiral components in a concentrated mixture can also phase separate into homochiral fluid domains, and the isomerization kinetics can then perform a symmetry breaking into one thermodynamical stable homochiral system. Glyceraldehyde has a sufficient chiral discrimination to perform such a symmetry breaking.

The requirement of a high concentration of the chiral reactant(s) in an aqueous solution in order to perform and *maintain* homochirality; the appearance of phosphorylation of almost all carbohydrates in the central machinery of life; the basic ideas that the biochemistry and the glycolysis and gluconeogenesis contain the trace of the biochemical evolution, all point in the direction of that homochirality was obtained just after- or at a phosphorylation of the very first products of the formose reaction, at high concentrations of the reactants in phosphate rich compartments in submarine hydrothermal vents.

A racemic solution of D,L-glyceraldehyde-3-phosphate could be the template for obtaining homochiral D-glyceraldehyde-3-phosphate(aq) as well as L-amino acids.

I. ORIGIN OF LIFE AND THE ENVIRONMENT AT EARTH 4 BILLION YEAR AGO

All carbohydrates and derivatives of carbohydrates in biosystems are D-configurations and all amino acids and derivatives of are L-configurations. This notation refers to the stereo specific configuration of ligands at the carbon atom next to the terminal -CH_2OH for carbohydrates and to the configuration at the (α-) carbon atom with a carboxylic acid and an amine group attached. The homochirality is crucial for the coupled biochemical reactions in living systems. The origin of this specific order could may be have been evolved during a purification of simple biosystems, but a general assumption is that the homochirality of carbohydrates and amino acids have been present at the origin of life on Earth.

The oldest undisputed evidence of life on Earth is 3 Gyr (billion years) old [10]. It is fossilized bacteria, but older findings in rocks dated to about 3.5 Gyr [1] and 3.8 Gyr [48] may also show sign of the presence of bacteria. From a biochemical and biological point of view a bacteria is, however, a very complex and highly ordered system and thus the purification of bio-molecules and the self-organisation into complex biological control systems must necessarily have appeared significantly earlier than the appearance of bacteria. The Earth is 4.56 Gyr old, and if life has originated on Earth, it has taken place in the time interval from the creation of the Earth and to the appearance of the first geological sign of life. To determine possible physicochemical models for the origin of homochirality and the origin of life it is thus necessary to know the physicochemical environment at this early time of the Earth's history.

Bacteria as well as all other biosystems are from a physicochemical point of view soft condensed matter, and one more fact is worth noticing. All the elementary bio-assembly of molecules are dehydrations (DNA,RNA,proteins, starch,fats) with release of one water molecule per polymerization step, with the consequence that H_2O must have been present at the self-organization of the complex biosystems at the origin of life. A part of the water in the oceans comes from meteors that contained ice and hit the Earth at the Late heavy bombardment 3.8–4 Gyr ago [16, 72]. But the solid mantle of the Earth also contains water [30] and there is evidence of oceans on Earth 4.4–4.3 Gyr long before the late heavy bombardment [49, 81]. So a natural conclusion from these simple facts is that life with homochirality of carbohydrates and amino acids started as self-organization in a fluid containing water as one of its components.

Another fact worth noticing is that all known biosystems only exist at moderate temperatures, $T \leq 150^{\circ}$C, and there are at least two reasons that life hardly could have started just after the

creation of the Earth. First the temperature just after the condensation of the materials, which has led to the creation of the planet(s) was too high for living systems with membranes. Secondly the Earth is believed to have been involved in a collision 30–100 Myr (million years) just after its creation, a collision which created the Moon [14, 15]. This collision and the creation of the Moon had drastic consequences not only for the temperature, but also for the condition at the surface of the Earth [82]. The moon was at the start much closer to the Earth and at that time a day was significant shorter. This is due to the tide waves which have pushed the Moon out to its present orbit and at the same time have damped the rotations of the Earth and the Moon. If Earth had oceans ≈ 4.3–4.4 Gyr ago there must have been very strong tide waves and a turbulent atmosphere. The extrapolation back to times just after the Moon's creation is difficult [7, 33], but all extrapolations indicate much stronger tide waves for $\approx$ 4–4.4 Gyr at the origin of life.

From a physicochemical point of view biosystems are non-equilibrium and open systems, and the refined self-organization must necessarily have taken place in an environment with high concentrations of the basic constituents and a smooth and continuous input over very long period of time of the constituents for the biosystems, which primarily are carbohydrates and amino acids. It is hard to imagine that this could have taken place in a turbulent atmosphere- or in the oceans of the Earth. But the right conditions could have been present at- or below the hydrothermal vents [64, 68].

Before analyzing the role of carbohydrates at the origin of homochirality in biosystems we need to summarize the appearance of carbohydrides and amino acids at the origin of life and their physicochemical behavior (Section 2), as well as the basic physicochemical-kinetics mechanism for obtaining and *maintaining* homochirality (Section 3).

II. AQUEOUS SOLUTIONS OF AMINO ACIDS AND CARBOHYDRATES

A. Prebiotic appearance of amino acids and carbohydrates on Earth

1. Amino acids

There are at least three sources to prebiotic amino acids: by lightning in the atmosphere, injected into the Earth from the interstellar space, or already present in the Earth's upper mantle or synthesized from the released materials at or below the vents.

The spectacular experiment(s) performed by [46] and [47] demonstrated that more than 20 dif-

ferent amino acids were produced by sparks ignition in a prebiotic atmosphere of water, methane, ammonia and hydrogen. In the original experiments Miller focused on amino acids, but later spark experiments demonstrated that also formaldehyde and other carbohydrates are synthesised [67]. So the lightning in this prebiotic atmosphere produced both the basic constituents in Earth's biological systems.

The carbonaceous meteorites contain amino acids and in a non-racemic proportion with a very small excess of L-amino acids. The composition of these meteorites is considered to be close to that of the solar nebula from which the solar system condensed. The two most prominent examples are the Murchison meteorite [5, 20] and the Murray meteorite [18]. From these meteorites compositional data and from an estimated impact rate of prebiotic meteorites, [56] suggested that meteorites might have contributed significantly by of the order up to 10^8 kg per year to amino acids on Earth at the time where the life could had started. But even with this injection of amino acids the concentration in the oceans must have been low. Earth's oceans now contain of the order 1.4×10^{21} kg H_2O. The simple amino acid glycine is observed in the interstellar space [39], and the interstellar space could have seeded the Earth with organic molecules at the origin of life [28].

Amino acids can also be synthesized at conditions similar to the physicochemical conditions at or below the hydrothermal vents [2, 31, 73]. If the materials from which the solar system has condensed contained the same materials as the carbonaceous meteorites, there might have been a substantial amount of amino acids in the upper part of the mantle of the Earth at the origin of life.

2. Carbohydrates

The formose reaction is the spontaneous condensation of formaldehyde into carbohydrides, which was discovered already in 1861 [13], and the aldol-like mechanism is described by [11]. The two first steps are

which gives a racemic mixture of D-glyceraldehyde and L-glyceraldehyde. Formaldehyde, CH_2O, is the simplest carbohydrate and it can be synthesized from carbon dioxide. It is produced in a reducing atmosphere containing CO_2 and methane [67]. The condensation into carbohydrides is catalyzed by not only amino acids [80]; but also by naturally aluminosilicates occurring at the alkaline hydrothermal vents [23, 44].

Simple carbohydrates are also detected in the interstellar space [28] which must have seeded the Earth, not only by amino acids but also with carbohydrates, and simple carbohydrates (formalde-





hyde) might also have appeared in the upper part of Earth's mantel at the origin of life.

The formose reaction is well known and is believed to be the basic synthesis of bio-carbohydrates and related bio-organic molecules [23, 32, 35, 41, 78].

B. Stability and isomerization kinetics of carbohydrates and amino acids

Both carbohydrates and amino acids are soluble in water due to the hydrophilic -OH groups in carbohydrates and to the hydrophilic -NH_2 and -COOH in amino acids which performs hydrogen bindings with water molecules. Both carbohydrates and amino acids are stable, but configurational unstable in the sense that they, by an isomerization kinetics, change configuration at an asymmetric carbon atom (i.e. an carbon atom with chemical bounds to four different atoms)

1. Carbohydrates

The kinetics which changes the configuration at a chiral center in a carbohydrate is a "keto-enol" kinetics. The simplest example is the kinetics for D-glyceraldehyde (GLA) in equilibrium with L-glyceraldehyde

Glyceraldehyde has only one asymmetric carbon atom and the keto-enol kinetic converts the D-configuration to the mirror L-configuration. The hexoses glucose and fructose have four and three asymmetric carbon atoms, respectively. The initiator of the configurational change by the keto-enol kinetics is the oxo group at a carbon atom next to an asymmetric carbon atom, which at a hexose is either carbon atom No. 1 (e.g. glucose) or to carbon atom No. 2 (fruc-



FIG. 1: Keto-enol isomerization kinetics for the triose glyceraldehyde

tose). The keto-enol intermediate is a configuration with two double bounded carbon atoms, each with an $-$OH group, (see Figure 1). It establishes a transformation between different configurational arrangements at carbon atoms next to the oxo group, in the case of glucose between D-glucose, D-fructose and D-manose, but not between the enantiomers D-glucose and L-glucose. The D-configuration of a carbohydrate refers to the configuration of the asymmetric carbon atom next to the terminal group -CH_2OH, in the case of glucose to carbon atom No. 5, and *this is not involved in the keto-enol kinetics*. To obtain a transformation between a D-carbohydrate and the enantiomer L-carbohydrate by the keto-enol kinetics one must go to the triose, as demonstrated in Figure 1. L-hexoses do not occur naturally in biosystems, but must be synthesized in the laboratory. (E.g. the acetate derivative of L-glucose is found to stimulate insulin release, but it is produced synthetic [43]).

Aqueous solutions of carbohydrates are reported to be unstable [38]. Reid and Orgel observed that diluted solutions of carbohydrates under prebiotic conditions degrade within hours or days [60] and ribose decomposes even at low temperature and neutral pH (44 years at $0< T \leq 150\,^{\circ}$C) [42]. None of the experiments contain, however, any information about the quality of the water, used for the solutions. Even highly purified water contains organic materials [83] and it requires special laboratories and technique to avoid bacteria in a solution. Bacteria, or their enzymes will degrade carbohydrates at low concentrations, but not at high concentrations. Carbohydrates at high concentrations are known to be very stable and they are used for conservation of food. Honey, which can be characterized as a fructose rich-solution with $\approx 17\%$ water is very stable, and fossil honey was found in in southern Georgia in a tomb from the bronze age, 27th–25th centuries B.C. [40].

2. Amino acids

Amino acids undergo also an isomerization kinetics [4], but there are crucial differences between the keto-enol kinetics for glyceraldehyde, pentoses and hexoses on one hand and the isomerization kinetics for bio-amino acids on the other hand. The pentoses and hexoses maintain their D-configuration at the keto-enol kinetics and only the carbohydrates with one asymmetric carbon, glyceraldehyde, changes between the D- and L-configuration. But all the chiral bio-amino acids change the L-configuration to the D-configuration by their isomerization kinetics. This is because all bio-amino acids are α-amino acids, i. e. the D- and L- configurations refer to the asymmetric carbon atom with an amino and a carboxylic group attached.

The isomerization kinetics will drive a diluted solution of a homochiral L-amino acid toward a racemic solution [4]. Another difference between carbohydrates and amino acids is the values of the reaction constants for the isomerization kinetics. Amino acids change configurations within thousands of years [4] (but might racemize much faster in the presence of a dipeptide [8]). Glyceraldehyde changes its configuration within hours, but with competing reactions [21].

Although the amino acids racemize they are, however, very (constitutional) stable in solid phases as well as in solutions, and even at relative high temperatures (423 K) [17, 69]. So if the composition of the carbonaceous meteorites equals the composition of the materials that created the Earth it is actual possible that the mantle of the Earth has contained amino acids at the origin homochirality and life.

III. SPONTANEOUS SYMMETRY BREAKING IN CONCENTRATED RACEMIC SOLUTIONS

Pasteur has demonstrated that the chiral components in a racemic mixture can separate into homochiral crystals of each enantiomers. A racemic system can, however, perform a spontaneous symmetry breaking to *one* homochiral system. There are *two* necessary conditions for that this process takes place. The reaction must be exergonic, i,e. $\Delta_r G < 0$ and the system must have an isomerization kinetics between the two chiral configurations. Some amino acids may fulfill these two conditions, but in the case of carbohydrates only the triose glyceraldehyde and its derivatives, e.g. glyceraldehyde-3-phosphate fulfills it. This is due to the fact that whereas the isomerization kinetics for α-amino acids converts D- to L- and visa verse, the keto-enol kinetics for higher-order carbohydrates as mentioned does not change the stereo configuration at the carbon atom next to

the terminal group.

A. Exergonic reaction in a racemic mixture of chiral molecules with strong chiral discrimination

The first and crucial condition for obtaining not only a spontaneous symmetry breaking, but also for maintaining homochirality over long times is that there is a (exergonic) gain of Gibbs free energy by a separation into homochiral sub domains. For the reaction

$$D, L \rightarrow D + L, \tag{1}$$

where D,L is an uniform mixture of molecules with D- and L configurations which separates into *sub domains* of the D and L forms, the the gain in Gibbs free energy,

$$\Delta_r G = \Delta_r H - T\Delta_r S \tag{2}$$

by this phase separation is primarily determined by the gain in enthalpy, $\Delta_r H < 0$ by the separation. This is because the change in free energy from the entropy term by the separation, $-T\Delta_r S > 0$, is positive and of the order $RT\ln 2 \approx 2\,\mathrm{kJ/mole}$ for a racemic mixture of equal amount of molecules with D- and L configurations (obtained by approximating the D,L mixture with an ideal mixture with mixing entropy equal to $R[x\ln x + (1-x)\ln(1-x)]$ and with mole fraction $x = 1/2$ for the racemic mixture).

The gain in enthalpy is given by the strength of the chiral discrimination (cd). The existence of chiral discrimination was discovered by Louis Pasteur in 1848 [54]. It manifests itself in the crystallization from a racemic fluid mixture of enantiomers into crystals of homochiral molecules, as demonstrated by Pasteur. As is well known, it is due to the fact that for some molecules there is an enthalpy gain, $-\Delta_r H > 0$, by packing homochiral molecules together instead of in pairs of enantiomers. The enthalpy gain must be bigger than the temperature times the mixing entropy, which is of the order 2 kJ/mole at room temperature. For pure systems, $\Delta_r H$ in Eq. (2) for the change in enthalpy by the phase separation due to the chiral discrimination can be obtained from standard values as

$$\Delta_r H = \Delta_{\mathrm{cd}} H^{\ominus} = \Delta_{\mathrm{f}} H^{\ominus}(\mathrm{enantiomer}) - \Delta_{\mathrm{f}} H^{\ominus}(\mathrm{racemic}) \tag{3}$$

From a physicochemical point of view biosystems are, however, in an aqueous solution. Because the polymerization of amino acids, as well as of carbohydrides and Nucleic acids, are dehydrations with a release of one water molecule per covalent polymerization bond, the state where the

synthesis of prebiotic systems appear is, what in physical chemistry is referred to by (aq) for the aqueous solution. On the other hand, the concentration of the chiral molecules, $[D]$ and $[L]$, at prebiotic self-assembling must necessarily have been high, with only a small concentration of water molecules, for two simple reasons. First

$$\Delta_{\rm cd}H \approx 0; [D], [L] \ll 1 \tag{4}$$

in diluted solution because there is no aggregation of homochiral clusters, so the reaction is endergonic ($\Delta_r G \approx -T\Delta_r S > 0$), and secondly the isomerization kinetics (the second general condition) in a diluted aqueous solution drives the system toward a racemic composition, as is observed in diluted aqueous solutions of L-amino acids [4]. So even if one, in one way or the other, starts with a homochiral system, it will tend to a racemic state if the concentration is low.

TABLE I: Strength of chiral discrimination for different molecules.

Molecule	$\Delta_{\rm cd}H^{\ominus}$/kJ mol^{-1}
Glyceraldehyde(l)	– 11.7
Phenylalanine(c)	– 6.7
Isoleucine(c)	– 5.4
Lysine(c)	$\approx$ 0.
Valine(c)	$\approx$ 0.
Leucine(c)	3.1
Serine(c)	6.3
Proline(c)	16.6
Threonine(c)	16.9
Alanine(c)	18.9

Data taken from NIST Standard Reference Database Number 69 and references therein.

The difference between Pasteur's observation of the spontaneous separation of enantiomers from a racemic mixture and a corresponding spontaneous separation for a racemic mixture of carbohydrates or amino acids is that Pasteur's chiral constituents: D- and L-ammonium sodium tartrate are moderately soluble in water, but carbohydrates and amino acids are very soluble. But even if it is possible to obtain a separation of e.g. amino acids by a crystallization, biosystems are soft condensed matter and we need to find a mechanism which not only separates the two

constituents in the racemic composition, but also stabilizes homochirality over (very) long times in a water environment. Therefore a measure of the strength of the chiral discrimination should be obtained for a state with high concentration of the chiral molecules and where

$$\Delta_r H \approx \Delta_{\rm cd} H^{\ominus}(l), \tag{5}$$

and this should at least be bigger (i.e. $\Delta_r H < -2\,$kJ/mole) than the mixing entropy to ensure an exergonic process.

In general, physicochemical data for the strength of chiral discrimination, $\Delta_{\rm cd} H^{\ominus}(l)$ in the fluid state do not exist. Only for the simple triose glyceraldehyde (Gla) where $\Delta_{\rm cd} H^{\ominus}(l) = -11.7\,$kJ/mole [6], which is much more than what is needed to overcome the mixing entropy. For some amino acids there exist data for $\Delta_{\rm cd} H^{\ominus}(c)$ for the enthalpy difference between an enantiomer crystal and a racemic crystal. This strength is, however, weakened at melting due to less effective packing in the fluid state. A collection of available data is given in Table 1. As can be seen from this table a few central amino acids (Phenylalanine, Isoleucine) might have the potential to maintain a homochiral aqueous state (aq). The simple triose, glyceraldehyde has, however, an exceptionally and sufficiently strong chiral discrimination.

B. The isomerization kinetics

The second general condition for obtaining a spontaneous symmetry breaking in a racemic mixture is the presence of an isomerization kinetics between the two enantiomeric configurations. The isomerization kinetics of amino acids and carbohydrides are complex [4, 21, 50]. We shall simplify the reaction mechanism by bimolecular reactions

$$\begin{array}{ccccc} & E_{\rm DD} & & E_{\rm DL} & \\ \mathrm{D + D} & \rightleftharpoons & \mathrm{D + L} & \rightleftharpoons & \mathrm{L + L} \\ & E_{\rm DL} & & E_{\rm LL} & \end{array} \tag{6}$$

between the two chiral species. The activation energy, $E_{\rm DL}$, for DL-collisions in an exergonic reaction, which may convert a D-molecule into a L-molecule or *vice versa*, is, in a condensed racemic fluid, less than the corresponding activation energy, $E_{\rm DD} = E_{\rm LL}$, thus allowing a conversion of one of the molecules in the collisions. The inequality

$$E_{\rm DL} < E_{\rm DD} = E_{\rm LL} \tag{7}$$

accounts for the chiral discrimination with a lower potential energy in homochiral domains of one of the enantiomers corresponding to a sufficient strength of the chiral discrimination [3]. More specifically

$$E_{\mathrm{DL}} + \mid \Delta_r H(x) \mid = E_{\mathrm{DD}} = E_{\mathrm{LL}}, \tag{8}$$

where the gain in enthalpy, $\mid \Delta_r H(x) \mid$, by a conversion of the stereo configuration is a function of the *local* composition, $x(\mathbf{r})$ of the configurations at the position $\mathbf{r}$, and with a maximum (negative) value of $\mid \Delta_r H(x) \mid$ for a homochiral local composition ($x(\mathbf{r}) = 0$ or $x(\mathbf{r}) = 1$).

Primarily $\mid \Delta_r H(x) \mid$ is proportional with the excess number of homochiral neighbors by a change of a configuration. Consider a simple example: Let a molecule, e.g. in a L configuration be activated to a transition state (intramolecular) configuration. It will with a Boltzmann probability end in the configuration with lowest energy. A simple molecule has twelve nearest neighbours. Let e.g. five of them be in a D configuration, four of the neighbours in a L-configuration and two be indifferent water molecules. It corresponds to that the local environment before the molecule was activated was racemic with an equal amount of D and L configurations. But the activated molecule will most likely turn into a D configuration by which the system tends to a lower energy, but now with an excess of D configuration. Thus a strong chiral discrimination together with an isomerization kinetics will ensure a separation of a racemic mixture on a molecular level and tend to order the molecules in homochiral clusters and sub domains.

A racemic D,L-system tends to separate in homochiral domains of enantiomers if the enthalpy gain, due to the chiral discrimination, is bigger than the entropy of mixing, just as in the case of Pasteur's experiment, but now in a fluid state. Without an isomerization kinetics such a racemic mixture will separate into homochiral sub domains through molecular diffusion. An isomerization kinetics enhances the separation; but what is much more significant, the kinetics also ensures a break of symmetry as demonstrated by the simple example, and results in the dominance of one of the species at late times [74].

The break of symmetry is obtained by, what could be expressed as *self-stabilizing chance* [77]. The deviation from a racemic mixture is self-stabilizing because a homochiral clusters- or sub domains catalyse their own growth by the isomerization kinetics which mainly takes place in the interface, whereas the chiral discrimination inside the homochiral domains slows down the kinetics and the conversions of the configurations, which always are unfavorable. Still one needs to explain the observed break of symmetry since the kinetics seems only to enhance the separation but *it does not favour one of the chiral species*. The break of symmetry on a macroscopic scale

will appear when one of the homochiral domains encapsulates the other domain. In a *confined geometry* of e.g. a fluid in a bottle or a prebiotic racemic mixture in a solid chamber [45, 63], one of the homochiral domains will dominate when, by a chance, it percolates in the confined volume and encapsulates the other domain.



FIG. 2: Illustrative example of the topological mechanism for symmetry breaking and the dominance of one of the chiral species. The (rectangular) volume contains the two separated homochiral phases of the enantiomers. The L-rich domain is white and it percolates the volume, whereas the red D-rich domain in the droplet is encapsulated by the L-rich domain. An activated particle (green) in the interface between the two domains has a range of attraction (blue) to the nearest neighbours which together with the activated particle determine the strength of the chiral discrimination. Since the white area inside the blue circle is greater than the corresponding red area, there are an excess of L neighbours and the activated particle will end in the L-configuration with a Boltzmann probability. The white area will increase and the area of the red droplet decrease and the droplet will finally disappear.

A simple example (Figure 2) illustrates this mechanism. Consider a system consisting of two homochiral (macroscopic) phases and let e.g. the D form be the dominating (red area in Figure 2), but as a big droplet in the middle of a box surrounded by molecules with L configurations (white area). The D molecules in the droplet has a convex surface (and the L molecules around the droplet have correspondingly a concave surface). Let the green circle mark an activated molecule A^* which will tend to either a D configuration or a L configuration with a Boltzmann probability depending on the local strength of $\Delta_r H(x(\mathbf{r}))$. Due to the strong chiral discrimination $\Delta_r H(x(\mathbf{r}))$ is only negative in the interface and the isomerization kinetics is mainly active there. Even if there are more D molecules in the droplet than L molecules around the droplet, most neighbours to

an activated molecule in the interface are L due to the concave L-rich neighbourhood, and the activated molecules will most likely end in a L configuration by which the (big) droplet decreases. It is a matter of *chance* which of the two enantiomers that dominates and encapsulates the other domain. The domain-catalytic behaviour is *self-stabilizing*, and the dominance of a percolating domain is self-stabilizing as well, because its concave interface will ensure the dominance. The topological determined condition for symmetry breaking and the dominance of one of the species is illustrated in Figure 2 [for further details see 74].

Figure 3 shows the time evolution of chiral dominance in a fluid of particles with chiral discrimination, obtained by a realistic (molecular dynamics) computer simulation [74]. The kinetics are implemented at time $t = 0$ and the strength of the chiral discrimination is given by the difference in the two activation energies, E_{DL} and $E_{\mathrm{DD}} = E_{\mathrm{LL}}$. The dominance of a species is measured by the enantiomeric excess given by the difference in mole fraction, $x_{\mathrm{D}} - x_{\mathrm{L}}$. For $E_{\mathrm{DD}} - E_{\mathrm{DL}} = RT$ we do not obtain a break of symmetry even after very long times, and inspection of the (local) composition shows that the mixture remains in the racemic state. This is also to be expected since the mixing entropy is $RT \ln 2$. But for a stronger chiral discrimination one observes a break of symmetry and a dominance of one of the enantiomers at late times. The figure only shows four cases; but many independent simulations gave an equal dominance of the D- and L- systems, within the statistical uncertainties, as one shall demand for a racemic system with no preference for one of the enantiomeric configurations. The dominance appears already for a strength of $2RT$ (b in the figure), but this strength is not sufficient to kill the isomerization kinetics within the dominating domain, and the system ends in a chiral fluid, but with a small homogeneous concentration of the loosing species. (Only for an infinite strong chiral dominance of $\Delta_r H(cd) = \infty$ (d in the figure) does this system end in a pure homochiral fluid.) Two other fluid systems have been investigated [75, 76] in order to determine the sensitivity of the break of symmetry of the actual fluid systems, the conclusion being, that the isomerization kinetics which destabilizes homochirality in diluted solutions ensures homochirality at high concentrations, provided that the chiral discrimination is sufficiently strong.

The description of the spontaneous symmetry breaking is exact. It makes only use of the law of thermodynamics and that the kinetics is reversible. But it is formulated for an isomerization kinetics between the two enantiomeric configurations in a concentrated aqueous solution. There are other models for symmetry breaking and spontaneous homochirality in racemic systems. Some of these models differs by assuming that the low Gibbs free energy state is not a concentrated



FIG. 3: The enantiomeric excess, i.e the difference in mole fraction, $x_{\mathrm{D}} - x_{\mathrm{L}}$, as a function of time in a molecular dynamics system with isomerization kinetics and for different strength of chiral discrimination, accordingly to Eq. (8): (a) $\mid \Delta_{cd} H^{\ominus} \mid = RT$, (b) $2RT$, (c) $3RT$ and (d) ∞.

aqueous solution of the basic constituent, but e.g. the crystalline phase or adsorbed constituents at mineral surfaces [27]. The general condition cited above is of cause also valid for these models, but the reason for focusing on concentrated aqueous solutions is that it is not enough to obtain a homochirality by crystallization or through adsorption at a mineral surface. A bacterium is, from a physicochemical point of view, soft condensed matter containing chiral molecules including both the L-amino acids and simple D-carbohydrates in an aqueous solution [19, 70], and the nucleus for life must from a physicochemical point of view, have been an open driven system with a steady state input of the chiral units for the self-assembly. So the homochirality must be maintained in this fluid over a very long period of time at the emergence of a living system.

In other models for the origin of homochirality in biosystems the chiral purification is obtained through the consecutive kinetic equations for the polymerization of the complex organic molecules. In the Frank model [9, 22] the composition of the simple chiral units in the biopolymers is assumed to be racemic at the origin of the synthesis, and the overall homochirality is obtained by the kinetics during a period of time at the origin of life. The homochiral domains are here the polymers and the chiral discrimination is given by the Gibbs free energy difference between the formation of a polymer binding between two homochiral centre versus the corresponding energy of a polymer binding between units with L- and D- configurations. This asymmetry is expressed by

the rate constants for cross-inhibition. This model is not at all in conflict with the present description. In both cases it is the kinetics in the open driven system which ensures homochirality, either by the isomerization kinetics between monomers or by the kinetics of polymerization, and in both cases governed by the gain of Gibbs free energy by homochiral (inter- or intramolecular) contacts. In fact both symmetry breaking kinetics might have been present at the origin of homochirality. But whereas the Frank model is an appealing model for purification of a racemic mixture of amino acids, the model is less appealing in the case of homochirality in polymers of carbohydrates such as the complex stereo-specific macromolecules DNA and RNA with a homochiral backbone of D-ribose. Here a model which ensures homochirality of the polymer units before the synthesis of the macromolecule is more appealing.

IV. HOMOCHIRALITY OF CARBOHYDRATES AND AMINO ACIDS

The carbohydrates, amino acids and their derivatives in biosystems are all homochiral. Thus one needs to explain how a symmetry breaking appears for both constituents. No matter where the ordering has taken place it is likely that both constituents have been present. But it is in fact very simple to explain this apparent puzzle. The mechanism works primarily through the gain of Gibbs free energy produced by a homochiral clustering together with an isomerization kinetics, and works equally well for mixtures of carbohydrates and amino acids provided there is a sufficient gain in energy by clustering D-carbohydrate molecules together with L-amino acid molecules. Nowhere in the thermodynamic/kinetic description in Section 3 for symmetry breaking in a concentrated aqueous solution is it assumed that all the molecules should be of the same type, only that there is a gain in energy by ordering and an active isomerization kinetics in the mixture. It is only a matter of whether there is a sufficient gain of Gibbs free energy by clustering amino acids together with a carbohydrate and this is actually the case for some simple carbohydrates and some amino acids [12, 36, 57, 58, 71].

In [36, 71] the authors observed a stereo selective coupling between L-serine octamers and D-glyceraldehyde and D-glucose. Serine furthermore performs coupling with phosphoric acid and some transition-metal ions. Serine has a unique ability to perform ”magic-number” ionic clusters composed of eight acid units and with a stereo selective chiral discrimination [51].

L-isovaline acts as a asymmetric catalyse of the threose and erythose from condensation of glycolaldehyde [57]. Glycolaldehyde and racemic glyceraldehyde condensate to pentoses in non

racemic proportions in the presence of (some) dipeptides of L,L- and D,D-amino acids [58].

The syntheses of glyceraldehyde from formaldehyde and glycolaldehyde in the present of homochiral amino acids gives a non racemic yield [12]. In [12] some L-amino acids were observed to give an excess amount of D-glyceraldehyde, e.g. L-phenylalanine, whereas e.g. L-proline gives a significant excess of L-glyceraldehyde.

These asymmetries, obtained in the synthesis of carbohydrates in the presence of excess L-amino acids, must be a result of the chiral discrimination (energy binding) between the carbohydrates and the amino acids. According to the model described above such gain of Gibbs free energy together with the isomerization kinetics of glyceraldehyde is sufficient to obtain a homochiral solution of D-glyceraldehyde. But the available experimental data show that only some of the L-amino acids favor an ordering of D,L-glyceraldehyde to D-glyceraldehyde. For some amino acids their L-configurations leads to an excess of L-glyceraldehyde [12, 58], e.g. L-proline. These catalysis works both ways so an excess of L-glyceraldehyde enhances the isomerization growth of L-proline, and correspondingly D-glyceraldehyde must enhance the isomerization growth of D-proline. According to Table 1 proline has a chiral discrimination which favors a racemic proportion even in a concentrated solution. Thus if the origin of homochirality of carbohydrates appeared via a catalyze by an L-amino acid, one needs to clarify how the homochirality of some L-amino acids, e.g. L-proline, has been obtained together with D-carbohydrates.

The chiral discrimination between amino acids and glyceraldehyde do not favour one of the constituents. But there is, however a significant difference between the isomerization kinetics of a carbohydrate and an amino acid. Whereas a diluted solution of an amino acid racemizes within thousand of years [4] [see however 8], the corresponding racemization of e.g. a diluted solution of D-glyceraldehyde is significant faster but with competing reactions [21]. On one hand this may be taken to imply that it is only possible within days or years to observe an excess of chirality of carbohydrates, but not of amino acids in a mixture of the two constituents, and one the other hand, that homochirality is first obtained for the carbohydrate no matter what the excess of chirality of the two constituents at the start of the purification.

V. THE ROLE OF PHOSPHATE AT THE ORIGIN OF LIFE

So far one important fact has been ignored. The role of phosphate at the origin of homochirality and life. Although all derivatives of carbohydrates have D-configurations they appear almost

always as phosphate derivatives. The list is long: The central energy cycles, the glycolysis and gluconeogenesis, ATP, ADP, RNA, DNA and the most common anchor molecule in the membrane molecules. Phosphates did not, however, appear in high concentrations in the early oceans [26, 34], but they are widely distributed in many minerals. Phosphate rocks containing apatite ($Ca_{10}(PO_4)_6(OH,F,Cl)_2$), are widespread on Earth and there are rare natural examples of condensed phosphates (e.g. canaphite, $Ca_2Na_2PO_7.4H_2O$ [62]; Kanonerovite $MnNa_3P_3O_{10}.12H_2O$ [59]).

A simple example illustrates the role of phosphate at the origin of life. There exist two different forms of simple living organism, Archaea and Bacteria. The central metabolisms (glycolysis and gluconeogenesis) are common but they differ on other biosynthetic pathways [24, 25], and by having different membrane molecules. Archaea membrane-molecules consist of (ether) derivatives of sn-glycerol-1-phosphate, whereas the molecules in the membranes of Bacteria and Eucarya consist of (ester) derivatives of sn-glycerol-3-phosphate. The membrane molecules are now synthesized by enzymes which, however, have no common sequence, and an analysis of these enzymes in different Achaea and Bacteria indicates that there is no common enzyme-ancestor [37]. In a living organism the membrane building blocks are synthesized from e.g. glycerol with help of these stereospecifique enzymes [55]; but at the origin of life we must search for a reaction pathway of spontaneous self-assembling directly from the ingredients. There are two facts which together are notable.

All the elementary membrane molecules in a cell are homochiral, and all these molecules (in Archaea and Bacteria membrane of three carbon atoms) can be synthesized from glyceraldehyde or glycerol. These two facts are notable because one can obtain simple vesicles with membranes from amphiphiles with only one or two central carbon atoms and with only one hydrophobic chain in the membrane molecule [53]. But nature did it with three carbon atoms in the central chiral unit and with two hydrophobic chains linked to the amphiphilic head group; all which can be synthesized from glyceraldehyde (see Figure 4). If the spontaneous self-assembly of cell-membranes had originated from a pool of simple bio-molecules, why would the spontaneous synthesis not have chosen the simplest track and synthesized simpler membrane molecules than diglycerides, phosphoro-lipids and phosphoro-ethers? The homochirality of the biomembranes are crucial for their functions and the first chiral component in the formose reaction is glyceraldehyde from which D-glyceraldehyde-3-phosphate and all the different membrane molecules can be synthesized.



FIG. 4: Phosphorylation and reduction of glyceraldehyde. If phosphorylation precedes reduction one obtains the configuration found in Archaea; if the reduction precedes phosphorylation one obtains both the configurations in Archaea and Bacteria. (Notice that due to $IUPAC$ nomenclature rules the reduced product of D-glyceraldehyde-3-phosphate is sn-glycerol-1-phosphate.)

VI. THE ROLE OF CARBOHYDRATES AT THE ORIGIN OF HOMOCHIRALITY IN BIOSYSTEMS

The general opinion and the literature about the origin of homochirality favors amino acids as the initiator and catalyzer of homochirality [29]. But inspection of the experiments and the present analysis reveals that it is more likely D,L-glyceraldehyde or rather D,L-glyceraldehyde-3-phosphate. The catalysis of symmetry breaking by the chiral discrimination of carbohydrates and amino acids is neutral in the sense that it does not select one of the reactants, e.g. an amino acid, as the catalyzer and the other as the substrate. A differences in the reaction times of the isomerization kinetics in favor of carbohydrates implies, however, that a carbohydrate reaches the homochiral state faster, and thus will act as the catalyzing template for homochiral purification of the amino acids. But if it was a carbohydrate that first reached the homochiral state, there seems to be only one possibility, glyceraldehyde or more likely the derivative glyceraldehyde-3-phosphate.



FIG. 5: The aldol condensation of dihydroxy acetone phosphate (DHAP) with D-glyceraldehyde-3-phosphate (D-GLA-3P) to D-fructose-1,6-diphosphate. The asymmetric carbon atom from D-GLA-3P which, by the condensation gives the D configuration in D-fructose-1,6-diphosphate is marked with *.

This is so because the isomerization kinetics of the higher order carbohydrates do not change the configuration at the carbon atom next to the terminal group, and because the spontaneous reaction of D,L-glyceraldehyde into pentoses and hexoses results in a spectrum of carbohydrates without an enantiomeric excess [79]. These fact favors a phosphorylation before or at a symmetry breaking and with the homochirality of D-sugars and L-amino acids obtained in a concentrated aqueous solution of D,L-glyceraldehyde-3-phosphate.

If the formose reaction took place in a phosphate environment with a phosphorylation before the condensation it explains not only the diversity of the enantiomeric form of the membrane molecules but also the central step in the glycolysis and gluconeogenesis. In this step dihydroxyacetone phosphate (DHAP) condensates with D-glyceraldehyde-3-phosphate (D-GLA-3-P) to D-fructose-1,6-diphosphate with the release of one water molecule (Figure 5). DHAP is one of the molecules in the isomerization reaction of D,L-glyceraldehyde-3-phosphate and will appear together with D-glyceraldehyde-3-phosphate. This aldol condensation is very exergonic at standard concentrations (molar concentrations at pH=7) with a Gibbs free energy $\Delta G^{\ominus'}$ = -23.98 kJ/mole and the reactants will react spontaneously [52]. The reason why we obtain energy from the reverse reaction with hydrolysis of D-fructose-1,6-diphosphate in the glycolysis is that the concentration of D-fructose-1,6-diphosphate is usual quite low ($<$0.1 mM) by which the reverse reaction is entropy driven and exergonic.

VII. IN SUMMARY

A break of symmetry and the ordering of a racemic solution into one homochiral domain only requires a strong chiral discrimination and an isomerization kinetics. These two conditions can be present in a concentrated racemic solution. The requirement of a high concentration of the chiral reactant(s) in an aqueous solution in order to perform and *maintain* homochirality; the appearance of phosphorylation of almost all carbohydrates in the central machinery of life; the basic ideas that the biochemistry and the glycolysis and gluconeogenesis contains the trace of the biochemical evolution [61, 66], all point in the direction of that homochirality was obtained just after- or at a phosphorylation of the very first products of the formose reaction at high concentrations of the reactants in phosphate-rich compartments.

An alkaline submarine hydrothermal mound is known to be compartmentalized and the inorganic membranes would offer low water activity traps and sites not only necessary for the phosphorylations [65] but also for the domain catalyzed and self-stabilizing mechanism by which the homochirality of simple organic chiral molecules can be obtained and maintained over long periods of time.

Although there have been many experimental investigations of possible spontaneous ordering of racemic systems into homochiral systems there do, however, not exist such experiments for D,L-glyceraldehyde-3-phosphate(aq).

Acknowledgments

The author acknowledge useful discussions with Axel Hunding and Jeppe C Dyre. The centre for viscous liquid dynamics “Glass and Time” is sponsored by the Danish National Research Foundation (DNRF) grant No. DNRF61.

[1] Altermann W, Kazmierczak J (2003) Archean microfossils: a reappraisal of early life on Earth. Res Microbiol 154:611-617

[2] Amend JP, Shock EL (1998) Energetics of Amino Acid Synthesis in Hydrothermal Ecosystems. Science 281:1659-1662

[3] Atkins P, de Paula J (2002) Atkins Physical Chemistry. Seventh Ed.; Oxford University Press; Chapter 27.5.

[4] Bada JL (1972) Kinetics of Razimization of Amino Acids as a Function of pH. J. Am. Chem. Soc. 94:1371-1373

[5] Bada JL, Cronin JR, Ho M-S, Kvenvolde KA, Lawless JG, Miller SL, Oro J, Steinberg S (1983) On the reported optical activity of amino acids in the Murchison meteorite. Nature 301:494-496

[6] Baer E, Flehmig H-H (1969) Refutation of alleged differences in energy contents of optical isomers. Can. J. Biochem. 47:79-83. Concerning the physical chemical state (l) see footnote to Table I.

[7] Bills BG, Ray RD (1999) Lunar Orbital Evolution: A Synthesis of Recent Results. Geophys. Res. Lett. 26:3045-3048

[8] Boehm MF, Bada JL (1984) Racemization of aspartic acid and phenylalanine in the sweetener aspartame at 100 degrees C. Proc. Nat. Acad. Sci. 81:5263-5266

[9] Brandenburg A, Andersen AC, Höfner S, Nilsson M (2005) Homochiral growth through enantiomeric cross-inhibition. Origin Life Evol. Biosphere 35:225-241

[10] Brasier M, McLoughlin N, Green O, Wacey D (2006) A fresh look at the fossil evidence for early Achaean cellular life. Philosophical Transactions of the Royal Society B 361:887-902

[11] Breslow R (1959) On the Mechanism of the Formose Reaction. Tetrahedron Lett. 1:22-26

[12] Breslow R, Cheng Z-L (2010) L-amino acids catalyze the formation of an excess of D-Glyceraldehyde, and thus of other D sugars, under credible prebiotic conditions. Proc. Nat. Acad. Sci. 107:5723-5725

[13] Butlerow A (1861) Formation synthetique d'une substance sucree. Compt. Rend. Acad. Sci. 53:145-147

[14] Canup RM (2004) Simulations of a late lunar-forming impact. Icarus 168:433-456

[15] Canup RM, Asphaug E (2001) Origin of the Moon in a giant impact near the end of the Earth's formation. Nature 412:708-712

[16] Cohen BA, Swindle TD, Kring DA (2000) Support for the Lunar Cataclysm Hypothesis from Lunar Meteorite Impact Melt Ages. Science 290:1754-1755

[17] Conway D, Libby WF (1958) The Measurement of Very Slow Reaction Rates: Decarcoxylation of Alanine. J. Am. Chem. Soc. 80:1077-1084

[18] Cronin JR, Moore C (1971) Amino Acid Analysis of the Murchison, Murray, and Allende Carbonaceous Chondrites. Science 172:1327-1329

[19] Ellis RJ (2001) Macromolecular crowding: obvious but underappreciated. Trends Biochem. Sci.

26:597-604

[20] Engel MH, Nagy B (1982) Distribution and enantiomeric composition of amino acids in the Murchison meteorite. Nature 296:837-840

[21] Fedoroňko M, Königstein J (1969) Kinetics of Mutual Isomerization of Trioses and their Dehydration to Methylglyoxal, Coll. Czechoslov. Chem. Commun. 34:3881-3894

[22] Frank FC (1953) On spontaneous asymmetric catalysis. Biochim. Biophys. Acta 11:459-463.

[23] Gabel N W, Ponnamperuma C (1967) Model for Origin of Monosaccharides. Nature 216:453-455

[24] Genschel (2004) Coenzyme A Biosynthesis: Reconstruction of the Pathway in Archaea and an Evolutionary Scenario Based on Comparative Genomics. Mol. Biol. Evol. 21:1242-1251

[25] Graham DE, White RH (2002) Elucidation of methanogenic coenzyme biosyntheses: from spectroscopy to genomics. Nat. Prod. Rep. 19:133-147

[26] Hagan WJ Jr., Parker A, Steuerwald A, Hathaway M (2007) Phosphate Solubility and the Cyanate-Mediated Synthesis of Pyrophosphate. Orig. Life Evol. Biosph. 37:113-122

[27] Hazen RM, Sverjensky DA (2010) Mineral Surfaces, Geochemical Complexities, and the Origin of Life. Cold Spring Harbor Perspectives in Biology doi: 10.1101

[28] Herbst E, van Dishoeck EF (2009) Complex Organic Interstellar Molecules. Annu. Rev. Astron. Astrophys. 47:427-480

[29] Higgs PG (2009) A Thermodynamic Basis for Prebiotic Amino Acid Synthesis and the Nature of the First Genetic Code. Astrobiology 9:483-490

[30] Hirschmann M, Kohlstedt D (2012) Water in Earth's mantle. Phys. today 65(3):40-45

[31] Huber C, Wächtershäuser G (2006) α-Hydroxy and α-Amino Acids Under Possible Hadean, Volcanic Origin-of-Life Conditions. Science 314:630-632

[32] Jalbout AF, Abrell L, Adamowicz L, Polt R, Apponi AJ, Ziurys LM (2007) Sugar synthesis from a gas-phase formose reaction. Astrobiol. 7:433-442

[33] Kagan B A (1997) Earth-Moon tidal evolution: model results and observational evidence. Prog. Oceanog. 40:109-124

[34] Keefe AD, Miller SL (1995) Are polyphosphates or phosphate esters prebiotic reagents? J. Mol. Evol. 41:693-702

[35] Kim H-J, Ricardo A, Illangkoon HI, Kim MJ, Carrigan MA, Frye F, Benner SA (2011) Synthesis of Carbohydrates in Mineral-Guided Prebiotic Cycles. J. Am. Chem. Soc. 133:9457-9468

[36] Kock KJ, Gozzo FC, Nanita SC, Takats Z, Eberlin MN, Cooks RG (2002) Chiral Transmission be-

tween Amino Acids: Chirally Selective Amino Acid Substitution in the Serine Octamer as a Possible Step in Homochirogenesis. Angew. Chem. Int. Ed. 41:1721-1724

[37] Koga Y, Kyuragi T, Sone N (1998) Did archeal and bacterial cells arise independently from noncellular precusors?. A hypothesis stating that the advent of membrane phospholipid with enantiomeric glycerophosphate backbones causes their separation of the two lines of descent. J. Mol. Evol. 46:54-63

[38] Kopetzki D, Antonietti M (2011) Hydrothermal formose reaction. New. J. Chem. 35:1787-1794

[39] Kuan Y-J, Charnley SB, Huang H-C, Tseng W-L, Kisiel Z (2003) Interstellar Glycine. Astrophys. J. 593:848-867

[40] Kvavadze E, Gambashidze I, Mindiashvili G, Gogochuri G (2007) The first find in southern Georgia of fossil honey from the Bronze Age, based on palynological data. Veget. Hist. Archaeobot. 16:399-404

[41] Lambert JB, Gurusamy-Thangavelu SA, Ma K (2010) The silica-mediated formose reaction: bottom-up synthesis of sugar silicates. Science 327:984-986

[42] Larralde R, Robertson MP, Miller SL (1995) Rates of decomposition of ribose and other sugars: Implications for chemical evolution. Proc. Natl. Acad. Sci. USA 92:8158-8160

[43] Malaisse WJ (1998) The riddle of L-Glucose pentaacetate insolinotropic action. Int. J. Mol. Med. 2:383-388

[44] Martin W, Baross J, Kelley D, Russell MJ (2008) Hydrothermal vents and the origin of life. Nature Reviews Microbiology 6:806-814

[45] Martin M, Russell MJ (2003) On the origins of cells: a hypothesis for the evolutionary transitions from abiotic geochemistry to chemoautotrophic prokaryotes, and from prokaryotes to nucleated cells. Phil. Trans. R. Soc. London Ser. B 358:59-85

[46] Miller SL (1953) Production of Amino Acids Under Possible Primitive Earth Conditions. Science 117:528- 529

[47] Miller SL, Urey HC (1959) Science 130:245-251

[48] Mojzsis SJ, Arrhenius G, McKeegan KD, Harrison TM, Nutman AP, Friend CRL (1996) Evidence for life on Earth before 3.8 Ga. Nature 384: 55-59

[49] Mojzsis SJ, Harrison TM, Pidgeon RT (2001) Oxygen-isotope evidence from ancient zircons for liquid water at the Earth's surface 4,300 Myr ago. Nature 409:178-181

[50] Nagorski RW, Richard JP (2001) Mechanistic Imperatives for Aldose-Ketose Isomerization in Water: Specific, General Base- and Metal Ion-Catalyzed Isomerization of glyceraldehyde with Proton and Hydride Transfer, J. Am. Chem. Soc. 123:794-802

[51] Nanita SC, Cooks RG (2006) Serine Octamers: Cluster Formation, Reactions, and Implications for Biomolecule Homochirality. Angew. Chem. Int. Ed. 45:554-569

[52] Nelson DL, Cox, MM (2010) Lehninger Principles of Biochemistry. Fifth Ed.,W. H. Freeman. Chapter 14.

[53] Ourisson G, Nakatani Y (1999) Origins of cellular life: molecular foundations and new approaches. Tetrahedron 55:3183-3190

[54] Pasteur ML (1848,1850) Recherches sur les Propriétés Spécifiques des deux Acides qui composent lÁcide Racémique. C. R. Acad. Sci. 26:535; Ann. Chim. Phys. 28:56-99

[55] Peretó J, Loópez-Garcia P, Moreira, D (2004) Ancestral lipid biosynthesis and early membrane evolution. Treds. Biol. Sci. 29:469-477

[56] Pierazzo E, Chyba CF (1999) Amino acid survival in large cometary impacts. Meteoritics Planet. Sci. 34:909-918

[57] Pizzarello S, Weber AL (2004) Prebiotic Amino Acids as Asymmetric Catalysts. Science 303:1151

[58] Pizzarello S, Weber AL (2010) Stereoselective Syntheses of Pentose Sugars Under Realistic Prebiotic Conditions". Origin Life Evol. Biosphere 40:3-10

[59] Popova VI, Popov VA, Sokolova EV, Ferraris G, Chukanov NV (2002) Kanonerovite, $MnNa_3P_3O_{10}.12H_2O$, first triphosphate mineral (Kazennitsa pegmatite, Middle Urals, Russia). N Jb Miner. Mh. Jg. 3:117-127

[60] Reid C, Orgel LE (1967) Synthesis of Sugar in Potentially Prebiotic Conditions. Nature 216:455

[61] Romano AH, Conway, T (1996) Evolution of carbohydrate metabolic pathways. Res. Microbiol. 147:448-455.

[62] Rouse RC, Peacor DR, Freed RL (1988) Pyrophosphate groups in the structure of canaphite, $Ca_2Na_2PO_7.4H_2O$; the first occurrence of a condensed phosphate as a mineral. American Mineralogist 73:168-171

[63] Russell MJ, Martin M (2004) The rocky roots of the acetyl COA pathway. Trends Biochem. Sci. 29:358-369

[64] Russell MJ, Hall AJ, Martin W (2010) Serpentinization as a source of energy at the origin of life. Geobiology 8:355-371

[65] Russell MJ, Nitschke W., Branscomp E (2013) The inevitable journey to being. Phil. Trans. R. Soc. Lond. B Biol. Sci. (in press)

[66] Say RF, Fuchs G (2010) Fructose 1,6-bisphosphate aldolase/phosphatase may be an ancestral gluco-

neogenic enzyme. Nature 464:1077-1081.

[67] Schlesinger G, Miller SL (1983) Prebiotic Synthesis in Atmospheres Containing CH_4, CO, and CO_2. J. Mol. Evol. 19:383-390

[68] Shock EL (1992) Chemical environments of submarine hydrothermal systems. Origins of Life and the Evolution of the Biosphere 22:67-107

[69] Snider MJ, Wolfenden R (2000) The Rate of Spontaneous Decarboxylation of Amino Acids. J. Am. Chem. Soc. 122:11507-11508

[70] Spitzer J, Poolman B (2009) The Role of Biomacromolecular Crowding: Ionic Strength, and Physicochemical Gradients in the Complexities of Life's Emergence. Microbiol. Mol. Biol. Rev. 73:371-388

[71] Takats Z, Nanita SC, Cooks RG (2003) Serine Octamer Reactions: Indicators of Prebiotic Relevance. Angew. Chem. Int. Ed. 42:3521-3523

[72] Tera F, Papanastassiou DA, Wasserburg GJ (1974) Isotopic evidence for a terminal lunar cataclysm. Earth Planet. Sci. Lett. 22:1-21

[73] Imai E, Honda H, Hatori K, Brack A, Matsuno K (1999) Elongation of Oligopeptides in a Simulated Submarine Hydrothermal System. Science 283:831-833

[74] Toxvaerd S (2000) Molecular Dynamics Simulations of Isomerization Kinetics in Condensed Fluids. Phys. Rev. Lett. 85:4747-4750

[75] Toxvaerd S (2004a) Droplet Formation in a Ternary-Fluid Mixture: Spontaneous Emulsion and Micelle Formation, J. Phys. Chem. 108:8641-8645

[76] Toxvaerd S (2004b) Domain catalyzed chemical reactions: A molecular dynamics simulation of isomerization kinetics. J. Chem. Phys. 120:6094-6099

[77] Toxvaerd S (2005) Homochirality in Bio-Organic Systems and glyceraldehyde in the Formose Reaction, J. Biol. Phys. 31:599-606

[78] Washington J (2000) The possible Role of Volcanic Aquifers in Prebiologic Genesis of Organic Compounds and RNA. Origin Life Evol. Biosphere 30:53-79

[79] Weber AL (1987) The triose model: glyceraldehyde as a source of energy and monomers for prebiotic condensation reactions. Origin Life 17:107-119

[80] Weber AL (2001) The Sugar Model: Catalysis by Amines and Amino Acid Products. Origin Life Evol. Biosphere 31:71-86

[81] Wilde SA, Valley JW, Peck WH, Graham C M (2001) Evidence from detrital zircons for the existence of continental crust and oceans on the Earth 4.4 Gyr ago. Nature 409:175-178

[82] Zahnle K, Schaefer L, Fegley B (2010) Earth's Earliest Atmospheres. Cold Spring Harb. Perspect. Biol. doi 10.1101

[83] see www.elgalabwater.com/water-quality-en.html